

Gravitoelectromagnetism and Dark Energy in Superconductors

Clovis Jacinto de Matos*

February 4, 2008

Abstract

A gravitomagnetic analogue of the London moment in superconductors can explain the anomalous Cooper pair mass excess reported by Janet Tate. Ultimately the gravitomagnetic London moment is attributed to the breaking of the principle of general covariance in superconductors. This naturally implies non-conservation of classical energy-momentum. Possible relation with the manifestation of dark energy in superconductors is questioned.

Keywords: Principle of General Covariance; Gravitomagnetism; Superconductivity; Dark energy.

1 Introduction

In 1989 Tate et *al.*[1] [2] reported a Cooper pair mass excess in Niobium of 84 parts per million greater than twice the free electron mass (m_e), whereas a theoretical calculation based on the theory of relativity predicts a value 8 parts per million less than $2m_e$. This disagreement between theory and experiment has not been resolved so far [3]–[6]. Conjecturing an additional gravitomagnetic term in the Cooper pairs canonical momentum accounts for Tate’s observations. This naturally leads to the conjecture that Tate’s excess of mass is not real but instead a rotating superconductor would simply exhibit a gravitomagnetic analogue to the well known magnetic London moment. However the magnitude of this conjectured gravitomagnetic field would be 10

*ESA-HQ, European Space Agency, 8-10 rue Mario Nikis, 75015 Paris, France, e-mail: Clovis.de.Matos@esa.int

orders of magnitude higher than Earth natural gravitomagnetic field (about 10^{-14} Rad/s).

The electromagnetic properties of superconductors can be explained through the breaking of gauge symmetry and consequently through a massive photon in the superconductive material. A similar mechanism for gravitation involving the breaking of the Principle of General Covariance (PGC) in superconductors would lead to a set of Proca type equations for gravitoelectromagnetism, with an associated massive spin one boson to convey the gravitoelectromagnetic interaction. Requiring that the PGC is recovered from this set of equations in the case of normal matter; we find back for the case of physical systems made simultaneously of coherent and normal matter, like superconductors, the anomalously high gravitomagnetic London moment conjectured from Tate's experiment. Ultimately It appears that Tate's measurements can be expressed in terms of the ratio between the Copper pair mass density, ρ_m^* , and the superconductor's bulk density, ρ_m .

$$\frac{m^* - m}{m} \simeq \frac{\rho_m^*}{\rho_m} \quad (1)$$

Where m^* and m are respectively the experimentally measured and the theoretically predicted Cooper pair mass.

The breaking of the PGC, implies that energy-momentum would not be conserved in superconductors. Could this be related with some still unknown properties of dark energy? The investigation of the physical nature of dark energy in quantum materials is a fascinating possibility, which Beck and Mackey[7] are already exploring for the case of Josephson junctions.

2 Gravitomagnetic London Moment

Tate et *al.* used a SQUID to measure the magnetic field generated by the rotation ($\omega = 2\pi\nu[\text{Rad/s}]$) of a thin (on the order of the London penetration depth) Niobium superconductive ring, also called the London moment. Following Ginzburg-Landau theory of superconductivity the total magnetic flux (including the Cooper pairs current density) cancels at regular frequency intervals $\Delta\nu[s^{-1}]$.

$$\frac{\hbar}{m^*} = 2S\Delta\nu \quad (2)$$

Where S is the area bounded by the Niobium ring. Based on Eq. (2) Tate estimated the Cooper pair's mass, $m^* = 1.82203 \times 10^{-30}[\text{Kg}]$, which is higher than the theoretically expected $m = 1.82186 \times 10^{-30}[\text{Kg}]$, including relativistic corrections[8].

In order to assess Tate's experiment, following DeWitt and Ross, a gravitomagnetic term is added to the Cooper pair's canonical momentum[9] [10], $\vec{p}_s$.

$$\vec{p}_s = m\vec{v}_s + e\vec{A} + m\vec{A}_g \quad (3)$$

where e is the Cooper pair's electric charge, $\vec{v}_s$ is the Cooper pair's velocity, $\vec{A}$ and $\vec{A}_g$ are respectively the magnetic and gravitomagnetic vector potentials.

Using again Ginzburg-Landau theory we find that the zero-flux condition, Eq. (2), depends now also on a gravitomagnetic component[11] [12].

$$\frac{\hbar}{m} = 2S\left(\Delta\nu + \frac{\Delta B_g}{4\pi}\right) \quad (4)$$

Doing Eq.(2) into Eq. (4) we find the gravitomagnetic variation needed to account for Tate's anomalous excess of mass[13] is: $\Delta B_g = 1.65 \times 10^{-5} [Rad/s]$.

Subtracting the classical Cooper pair's canonical momentum: $\vec{p}_s^* = m^*\vec{v}_s + e\vec{A}$, from the generalised one, Eq. (3), and taking the rotational, we obtain the gravitomagnetic analogue of the London moment in superconductors, which we call the "Gravitomagnetic London Moment"[14].

$$\vec{B}_g = \frac{m^* - m}{m} 2\vec{\omega} = \frac{\Delta m}{m} 2\vec{\omega} \quad (5)$$

Therefore we conclude that the Cooper pair's mass does not increase but instead when a superconductor is set rotating it generates simultaneously a (homogeneous) magnetic field (London Moment) together with an (homogeneous) gravitomagnetic field (Gravitomagnetic London Moment). If the latest phenomenon is neglected it is naturally interpreted, as Tate did, as being an anomalous excess of mass of the Cooper pairs.

The gravitomagnetic field generated by the rotating Niobium ring in Tate's experiment would then be:

$$\vec{B}_g = 1.84 \times 10^{-4} \vec{\omega}, \quad (6)$$

which is very large even for small angular velocities compared to classical astronomical sources, but cannot be ruled out based on the experimental results achieved so far[15]. However how can we explain this conjecture? We will see that the answer passes through the investigation of the validity of the Principle of General Covariance for superconductors.

3 Spontaneous Breaking of Gauge Invariance in Superconductors

Superconductor's properties (zero resistivity, Meissner effect, London moment, flux quantization, Josephson effect etc...) can be understood likewise

spontaneous breaking of electromagnetic gauge invariance when the material is in the superconductive phase[16] [17]. In field theory, this symmetry breaking leads to massive photons via the Higgs mechanism. In this case the Maxwell equations transform to the so-called Maxwell-Proca equations, which are given by:

$$\nabla \vec{E} = \frac{\rho^*}{\epsilon_0} - \frac{1}{\lambda_\gamma^2} \phi, \quad (7)$$

$$\nabla \vec{B} = 0, \quad (8)$$

$$\nabla \times \vec{E} = -\dot{\vec{B}}, \quad (9)$$

$$\nabla \times \vec{B} = \mu_0 \rho^* \vec{v}_s + \frac{1}{c^2} \dot{\vec{E}} - \frac{1}{\lambda_\gamma^2} \vec{A}, \quad (10)$$

Where $\vec{E}$ is the electric field, $\vec{B}$ is the magnetic field, ϵ_0 is the vacuum electric permittivity, $\mu_0 = 1/\epsilon_0 c^2$ is vacuum magnetic permeability, ϕ is the scalar electric potential, $\vec{A}$ is the magnetic vector potential, ρ^* is the Cooper pairs fluid electric density, $\vec{v}_s$ is the cooper pairs velocity, and $\lambda_\gamma = \hbar/m_\gamma c$ is the photon's Compton wavelength, which is equal to the London penetration depth $\lambda_L = \sqrt{\frac{m}{\mu_0 \rho^* e}}$.

Taking the rotational of Eq. (10) and neglecting the term coming from the displacement current, we get the following equation for the magnetic field:

$$\nabla^2 \vec{B} + \frac{1}{\lambda_\gamma^2} \vec{B} = \frac{1}{\lambda_L^2} \frac{m}{e} 2\vec{\omega}. \quad (11)$$

Solving Eq. (11) for a one dimensional case, we obtain the Meissner effect and the London moment.

$$B = B_0 e^{-x/\lambda_\gamma} + 2\omega \frac{m}{e} \left(\frac{\lambda_\gamma}{\lambda_L} \right)^2 \quad (12)$$

Following the argument from Becker et al.[18] and London [19], the London moment is developed by a net current that is lagging behind the positive lattice matrix, The Cooper pair current density direction sign has to show in opposite direction than the angular velocity of the superconducting bulk. This is important as the London moment in all measurements, due to the negative charge of the Cooper pair, shows in the same direction as the angular velocity. Having $\lambda_\gamma = \lambda_L$ we finally get:

$$B = B_0 e^{-x/\lambda_L} - 2\omega \frac{m}{e} \quad (13)$$

4 Spontaneous Breaking of the Principle of General Covariance in Superconductors

General Relativity is founded on the *principle of equivalence*, which rests on the equality between the inertial and the gravitational mass of any physical system, and formulates that *at every space-time point in an arbitrary gravitational field it is possible to choose a "locally inertial coordinate system" such that, within a sufficiently small region of the point in question, the laws of nature take the same form as in unaccelerated Cartesian coordinate systems in the absence of gravity*. In other words, The inertial frames, that is, the "freely falling coordinate systems", are indeed determined by the local gravitational field, which arises from all the matter in the universe, far and near. However, once in an inertial frame, the laws of motion are completely unaffected by the presence of nearby masses, either gravitationally or in any other way.

Following Steven Weinberg, the *Principle of General Covariance* (PGC) is an alternative version of the principle of equivalence[20], which is very appropriate to investigate the field equations for electromagnetism and gravitation. It states that *a physical equation holds in a general gravitational field, if two conditions are met:*

1. The equation holds in the absence of gravitation; that is, it agrees with the laws of special relativity when the metric tensor $g_{\alpha\beta}$ equals the Minkowsky tensor $\eta_{\alpha\beta}$ and when the affine connection $\Gamma_{\beta\gamma}^{\alpha}$ vanishes.
2. The equation is generally covariant; that is, it preserves its form under a general coordinate transformation $x \rightarrow x'$.

It should be stressed that general covariance by itself is empty of physical content. The significance of the principle of general covariance lies in its statement about the effects of gravitation, that a physical equation by virtue of its general covariance will be true in a gravitational field if it is true in the absence of gravitation. The PGC is not an invariance principle, like the principle of Galilean or special relativity, but is instead a statement about the effects of gravitation, and about nothing else. In particular general covariance does not imply Lorentz invariance. Any physical principle such as the PGC, which takes the form of an invariance principle but whose content is actually limited to a restriction on the interaction of one particular field, is called a dynamic symmetry. As discussed above local gauge invariance, which governs the electromagnetic interaction is another important dynamical symmetry. We can actually say that the Principle of General Covariance

in general relativity is the analogous of the Principle of Gauge Invariance in electrodynamics.

Maxwell-Proca equations for electromagnetism Eqs.7-10, which apply in a superconductor, are not gauge invariant just as they are not generally covariant. If we assume that the PGC is spontaneously broken in a superconductor, like gauge invariance is, the weak field approximation of Einstein field equations would lead to the following set of Proca equations for gravito-electromagnetism, which contains a spin 1 massive boson, called graviphoton, to convey the gravitoelectromagnetic interaction [22] [23].

$$\nabla \vec{g} = -\frac{\rho_m^*}{\epsilon_{0g}} - \frac{1}{\lambda_g^2} \phi_g, \quad (14)$$

$$\nabla \vec{B}_g = 0, \quad (15)$$

$$\nabla \times \vec{g} = -\dot{\vec{B}}_g, \quad (16)$$

$$\nabla \times \vec{B}_g = -\mu_{0g} \rho_m^* \vec{v}_s + \frac{1}{c^2} \dot{\vec{g}} - \frac{1}{\lambda_g^2} \vec{A}_g, \quad (17)$$

Where $\vec{g}$ is the gravitational field, $\vec{B}_g$ is the gravitomagnetic field, $\epsilon_{0g} = 1/4\pi G$ is the vacuum gravitational permittivity, $\mu_{0g} = 4\pi G/c^2$ is vacuum gravitomagnetic permeability, ϕ_g is the scalar gravitational potential, $\vec{A}_g$ is the gravitomagnetic vector potential, ρ_m^* is the Cooper pairs mass density, $\vec{v}_s$ is the cooper pairs velocity, and $\lambda_g = \hbar/m_g c$ is the Compton wavelength of the graviphoton.

Taking the gradient of Eq. (14), and the rotational of Eq. (17), and solving the resulting differential equations for the one dimensional case we find respectively the form of the principle of equivalence and of the gravitomagnetic Larmor Theorem[24] in superconductive cavities.

$$\vec{g} = -\vec{a} \mu_{0g} \rho_m^* \lambda_g^2, \quad (18)$$

$$\vec{B}_g = 2\vec{\omega} \mu_{0g} \rho_m^* \lambda_g^2 \quad (19)$$

where for Eq. (19) we had to introduce Becker's argument that the Cooper pairs are lagging behind the lattice so that the current is flowing in the opposite direction of ω .

In order to find a phenomenological law for the graviphoton wavelength[25] we request that from Eq. (18) and (19) the PGC is restored in the case of normal matter. In that case the mass density reduces to the materials bulk density, ρ_m , no condensate phase is present within the material.

$$\vec{g} = -\vec{a} \mu_{0g} \rho_m \lambda_g^2, \quad (20)$$

$$\vec{B}_g = 2\omega \mu_{0g} \rho_m \lambda_g^2 \quad (21)$$

Since normal matter complies with the PGC, Eqs. (20) and (21) must reduce to:

$$\vec{g} = -\vec{a}, \quad (22)$$

$$\vec{B}_g = 2\vec{\omega} \quad (23)$$

Comparing Eq. (22) and (23) with Eq. (20) and (21) we find that the gravitphoton Compton wavelength must be inversely proportional to the local density of the bulk material mass:

$$\frac{1}{\lambda_g^2} = \mu_{0g}\rho_m \quad (24)$$

Doing Eq. (24) into Eq. (18) and (19) we get:

$$\vec{g} = -\vec{a}\frac{\rho_m^*}{\rho_m}, \quad (25)$$

$$\vec{B}_g = 2\vec{\omega}\frac{\rho_m^*}{\rho_m} \quad (26)$$

which clearly indicate a breaking of general covariance. Notice that in the case of Bose Einstein Condensates (BEC) we have only one single condensate phase in our material ($\rho_m^* = \rho_m$) implying that the PGC is no violated in BECs.

Comparing Eq. (26) with Eq. (5) and (6) we can explain the additional gravitomagnetic term in the Cooper pairs canonical momentum, in function of the mass density ratio of the coherent and normal phase.

$$9.2 \times 10^{-5} = \frac{m^* - m}{m} \simeq \frac{\rho_m^*}{\rho_m} = 3.9 \times 10^{-6} \quad (27)$$

The numerical values in Eq. (27) correspond to the case on Niobium in Tate's experimental conditions.

5 Dark Energy in Superconductors?

It is well known that breaking of general covariance leads to non-conservation of energy-momentum (in the covariant sense)[20]. Would that mean that in a superconductor we could observe some manifestation of dark energy?

Presently the physical nature of dark energy is unknown. What is clear is that various astronomical observations (supernovae, CMB fluctuations, large-scale structure) provide rather convincing evidence that around 73 percent of

the energy contents of the universe is a rather homogeneous form of energy, so-called "dark energy". A large number of theoretical models exist for dark energy, but an entirely convincing theoretical breakthrough has not yet been achieved. Popular models are based on quintessence fields, phantom fields, quintom fields, Born-Infeld quantum condensates, the Chaplygin gas, fields with non-standard kinetic terms, possible links between the cosmological constant and the graviton mass [21] to name just a few (see for example Refs. [27] and [28] for reviews). All of these approaches contain "new physics" in one way or another, though at different levels. However, it is clear that the number of possible dark energy models that are based on new physics is infinite. Only experiments will ultimately be able to confirm or refute the various theoretical constructs[26].

Beck is currently exploring the possibility that vacuum fluctuations, allowed by the uncertainty relation, in Josephson junction create dark energy. This is a priori the simplest explanation for dark energy. Assuming that the total vacuum energy density associated with zero-point fluctuations cannot exceed the presently measured dark energy density of the universe, Beck predicts an upper cutoff frequency of $\nu_c = (1.69 \pm 0.05) \times 10^{12} Hz$ for the measured frequency spectrum of the zero point fluctuations in the Josephson junction. The largest frequencies that have been reached in the experiments are of the same order of magnitude as ν_c and provide a lower bound on the dark energy density of the universe. If this is confirmed by future experiments how would that be related with the photon and graviphoton mass in superconductors? Where does the mass of these particles come from in the superconductive material? Finally Can we break general covariance in a superconductor without violating energy momentum conservation, if a superconductor contains a new form of energy?

6 Conclusion

The close analogy between the Principle of General Covariance and Gauge invariance, allows us to investigate the gravitoelectromagnetic properties of quantum materials in the framework of massive gravitoelectromagnetic Proca equations. We find that the breaking of the PGC in superconductors leads to a gravitomagnetic London moment and an associated additional gravitomagnetic term in the Cooper pairs canonical momentum, which can explain the anomalous excess of mass of Cooper pairs reported by Tate.

The breaking of the PGC in superconductors implies the non-equivalence between a rigid reference frame made with superconductive walls (superconductive cavity), being uniformly accelerated in a gravitational field free

region,

$$\vec{g} = -\vec{a}\left(1 + \frac{\rho_m^*}{\rho_m}\right), \quad (28)$$

$$\vec{B}_g = 2\vec{\omega}\left(1 + \frac{\rho_m^*}{\rho_m}\right) \quad (29)$$

and a classical rigid reference frame (made with normal matter) in a similar situation.

$$\vec{g} = -\vec{a}, \quad \vec{B}_g = 2\vec{\omega} \quad (30)$$

However breaking the principle of general covariance leads to a violation of the law of conservation of energy-momentum. It is not clear yet if this would be a sign for some manifestation of dark energy in superconductive materials. However it worths further investigation.

7 Acknowledgement

I am grateful to Prof. Orfeu Bertolami, Prof. John Moffat, Prof. Francis Everitt, Prof. Alan Kostelecky, for fruitful discussions and pertinent comments. I would like also to thank Prof. Raymond Chiao for encouragements and for being a rich source of inspiration to the present work. My profound gratitude also goes to Dr. Slava Turyshev and to all the organizers of the "From Quantum to Cosmos" conference for the fantastic forum of debates, they succeeded to establish during the conference.

References

- [1] J. Tate, B. Cabrera, S. B. Felch, J. T. Anderson, *Phys. Rev. Lett.* **62**, (8) 845 (1989).
- [2] J. Tate, B. Cabrera, S. B. Felch, J. T. Anderson, *Phys. Rev. B* **42**, (13) 7885 (1990).
- [3] M. Liu, *Phys. Rev. Lett.* **81**, (15) 3223 (1998).
- [4] Y. Jiang, M. Liu, *Phys. Rev. B* **63**, 184506 (2001).
- [5] H. Capellmann, *Europ. Phys. Jour. B* **25**, 25 (2002).
- [6] K. Capelle, E. K. U. Gross, *Phys. Rev. B* **59**, (10) (1999).
- [7] C. Beck, M. C. Mackey, *Phys. Lett. b* **605**, 295 (2005).

- [8] B. Cabrera, H. Gutfreund, W. A. Little, *Phys. Rev. B* **25**, (11) 6644 (1982).
- [9] B. S. DeWitt, *Phys. Rev. Lett.* **16**, 1092 (1966).
- [10] D. K. Ross, *Jour. Phys. A* **16**, 1092 (1983).
- [11] C. Kiefer, C. Weber, *Annalen Phys* **14**, 253 (2005).
- [12] O. Bertolami, M. Tajmar *Gravity Control and Possible Influence on Space Propulsion*, (ESA Contract Report CR(P)4365) (2002)
- [13] M. Tajmar, C. J. de Matos, *Physica C* **385**, 551 (2003).
- [14] M. Tajmar, C. J. de Matos, *Physica C* **420**, 56 (2005).
- [15] M. Tajmar, F. Plesescu, K. MArhold, C. J. de Matos, *gr-qc:0603033*, (2006).
- [16] L. H. Ryder, *Quantum Field Theory*, 2nd edn. (Cambridge University Press, 1996), p. 296.
- [17] S. Weinberg, *The Quantum Theory of Fields*, Vol 2, (Cambridge University Press, 1996),p. 332.
- [18] R. Becker, G. Heller, F. Sauter, *Z. Physic* **85**, 772 (1933).
- [19] F. London, *Superfluids*, (John Wiley and Sons, New York, 1950).
- [20] S. Weinberg, *Gravitation and Cosmology Principles and Applications of the General Theory of Relativity*, (John Wiley and Sons, New York, 1972) p.91, 111, 361.
- [21] M. Tajmar, Local cosmological Constant and the Dark Energy Coincidence Problem, to appear in *Class. Quant. Grav.*
- [22] J. Argyris, *Aust. J. Phys.* **50**, 879 (1997).
- [23] C. J. de Matos, M. Tajmar, *Physica C* **432**, 167 (2005).
- [24] B. Mashhoon, *Phys. Lett. A* **173**, 347 (1993).
- [25] M. Tajmar, C. J. de Matos, *gr-qc/0603032*, (2006).
- [26] C. Beck, *J.Phys.Conf.Ser.* **31**, 123 (2006).
- [27] P. J. E. Peebles, B. Ratra, *Rev. Mod. Phys.* **75**, 559 (2003)
- [28] T. Padmanabhan, *Phys. Rep.* **380**, 235 (2005)